arXiv:1302.4686v1 [math-ph] 19 Feb 2013

CATEGORIFICATION OF THE HEISENBERG ALGEBRA AND MACMAHON FUNCTION

NA WANG, ZHIXI WANG, KE WU, JIE YANG, AND ZIFENG YANG

Abstract. Starting from a one dimensional vector space, we construct a categorification $'\mathcal{H}$ of a deformed Heiserberg algebra $'H$ by Cautis and Licata's method. The Grothendieck ring of $'\mathcal{H}$ is $'H$. As an application, we discuss some related partition functions related to the MacMahon function of 3D Young diagram. We expect further applications of the results of this paper.

1. Introduction

Let H be the classical Heisenberg algebra of infinite rank, i.e., the algebra generated by the sequence $\{a_n\}_{n\in\mathbb{Z}}$ with the defining relations

$$a_na_m = a_ma_n + n\delta_{n+m,0}1, \quad n, m \in \mathbb{Z}. \tag{1}$$

The generators a_n for $n \in \mathbb{Z}$ can be realized from the fermions $\phi_n, \phi_n^*(n \in \mathbb{Z})$, which satisfy the following relations

$$[\phi_n, \phi_m]_+ = 0,\ [\phi_n^*, \phi_m^*] = 0,\ [\phi_n^*, \phi_m]_+ = \delta_{n+m,0}, \tag{2}$$

where we use the notation $[X, Y]_+ = XY + YX$. Let $\mathcal{A}$ be the Clifford algebra generated by the $\phi_n, \phi_n^*, n \in \mathbb{Z}$. Then an element of $\mathcal{A}$ can be written as a finite linear combination of monomials of the form

$$\phi_{m_1}\cdots\phi_{m_r}\phi_{n_1}^*\cdots\phi_{n_s}^*, \quad \text{where } m_1 < \cdots < m_r,\ n_1 < \cdots < n_s.$$

By introducing a variable k, we define the fermionic generating functions as the formal sums

$$\phi(k) = \sum_{n\in\mathbb{Z}+\frac{1}{2}} \phi_n k^{-n-\frac{1}{2}}, \quad \phi^*(k) = \sum_{n\in\mathbb{Z}+\frac{1}{2}} \phi_n^* k^{-n-\frac{1}{2}}.$$

Then the a_n can be realized by the following equation

$$\sum_{n\in\mathbb{Z}} a_n k^{-n-1} =: \phi(k)\phi^*(k) : \tag{3}$$

where :: is the normal order, see [MJD] for detail.

The integral form $H_{\mathbb{Z}}$ of Heisenberg algebra H using the vertex operators in terms of a_n is generated by $\{p_n, q_n\}_{n\in\mathbb{Z}}$ with relations

$$\begin{aligned} q_np_m &= \sum_{k\geq 0} p_{m-k}q_{n-k}, & (4)\\ q_nq_m &= q_mq_n, & (5)\\ p_np_m &= p_mp_n, & (6)\end{aligned}$$

in which $p_0 = q_0 = 1$ and $p_k = q_k = 0$ for $k < 0$, thus the summation in Equation (4) is finite. These generators $q_n,\ p_m$ are obtained as the homogeneous components in z of the halves of vertex operators

$$\sum_{m\geq 0} q_m z^{-m} = \exp(\sum_{m\geq 1} \frac{1}{m} a_m z^{-m}), \qquad \text{and} \qquad \sum_{m\geq 0} p_m z^m = \exp(\sum_{m\geq 1} \frac{1}{m} a_{-m} z^m),$$

which play a very important role in the conformal field theory and the theory of quantum algebras.

The authors use slightly different presentations of integral forms of Heisenberg algebra in literature. Khovanov used an integral form $H'_{\mathbb{Z}}$ of Heisenberg algebra in [K]. He introduced a calculus of planar diagrams for biadjoint functors and degenerate affine Hecke algebras, which led to an additive monoidal category whose Grothendieck ring contains the integral form $H'_{\mathbb{Z}}$ (and Khovanov conjectured that they are isomorphic).

We define the deformed integral form $'H_{\mathbb{Z}}$ of Heisenberg algebra as the algebra generated by $\{p_n, q_n\}_{n\in\mathbb{Z}}$ with relations

$$\begin{aligned} q_n p_m &= \sum_{k\geq 0}[k+1]p_{m-k}q_{n-k}, && (7)\\ q_n q_m &= q_m q_n, && (8)\\ p_n p_m &= p_m p_n, && (9)\end{aligned}$$

where $p_0 = q_0 = 1, \quad p_k = q_k = 0$ for $k < 0$, and the quantum integer $[k] = \frac{t^k-1}{t-1}$. We see that $'H_{\mathbb{Z}}$ is $H_{\mathbb{Z}}$ when $t = 0$.

It is important to lift the Heisenberg algebra (any form) to some categoric version in representation theory. Cautis and Licata [CL] categorified the Heisenberg algebras $\mathfrak{h}_\Gamma$ for some finite subgroup Γ of $SL_2(\mathbb{C})$, but the discussion in that paper clearly can not be applied to the case when Γ is the trivial group. We will give a categorification of the Heisenberg algebra $'H_{\mathbb{Z}}$ by the same techinique, and derive the MacMahon function from this categorification. The results of this paper are expected to be applied in the study of topics in the theory of quantum algebras which we are undergoing.

The authors are grateful to Morningside Center of Chinese Academy of Sciences for providing excellent research environment and financial support. This work is also partially supported by NSF grant 11031005 and grant KZ201210028032.

2. Categorification of the deformed Heisenberg algebra $'H$

Let $'H$ denote the Heisenberg algebra with a coefficient extension of $'H_{\mathbb{Z}}$ to some given field, and all the Grothendieck ring with the same coefficient extension.

Let $V = \Bbbk v$ be a 1-dimensional vector space over an algebraically closed $\Bbbk$ field of characteristic zero, $\Lambda^*(V)$ the exterior algebra of V with the basis B consisting of 1, v. We define a $\Bbbk$-linear map $tr: \ \Lambda^*(V) \to \Bbbk$ by setting

$$tr(v) = 1, \quad tr(1) = 0.$$

Let $B^\vee$ be the basis of $\Lambda^*(V)$ dual to B with respect to the non-degenerate bilinear form $\langle a, b\rangle := \mathrm{tr}(ab)$. Under this bilinear form, we see that $1^\vee = v, \ v^\vee = 1$.

We define a 2-category $'\mathcal{H}'$ as follows. The objects of $'\mathcal{H}'$ are integers. The 1-morphisms are generated by $P: \ n \to n+1$ and $Q: \ n+1 \to n$ by finite compositions and finite direct sums, here we use the same notation P or Q for different $n \in \mathbb{Z}$. We also denote by the same notation $\mathbf{1}$ the identity 1-morphisms on the objects. The 2-morphisms are planar diagrams modular relative to boundary isotopies. An upward oriented strand denotes the identity 2-morphism id : $P \to P$ while a downward oriented strand denotes the identity 2-morphism id : $Q \to Q$. Upward-oriented lines and downward-oriented lines carrying dots labeled by elements $b \in \Lambda^*(V)$ are other 2-morphisms of $'\mathcal{H}'$, for example,



$\in \mathrm{Hom}_{'\mathcal{H}'}(P, P)$,



$\in \mathrm{Hom}_{'\mathcal{H}'}(Q, Q)$,



$\in \mathrm{Hom}_{'\mathcal{H}'}(QP, PQ)$.

On planar diagrams, compositions of 2-morphisms are depicted upward from bottom to top. All these are similar to [CL], so are the local relations which the planar diagrams are requested to satisfy.

The local relations are the following. First, the dots can move freely along strands and through intersections, for example,



Second, collision of dots is controlled by the multiplication in the exterior algebra $\Lambda^*(V)$:



Third, dots on strands supercommute when they move past one another, for example:



Last, local relations contain the following:



(10)



(11)



(12)



(13)

We also define the degrees of the planar diagrams:



and the degree of a dot labeled by b equals the degree of b in the graded algebra $\Lambda^*(V)$. Thus for any two 1-morphisms A, B, the vector space of 2-morphisms $A \to B$ are graded, and the composition of

2-morphisms is compatible with grading. In addition, we define the shifted degrees (which is denoted by *sdeg* of the planar diagrams:

$sdeg$  $= 0$;

$sdeg$  $= -1/2$, $sdeg$  $= -1/2$;

$sdeg$  $= 1/2$, $sdeg$  $= 1/2$;

and the shifted degree $sdeg(b)$ of a dot labeled by $b \in \Lambda^*(V)$ is the same as $deg(b)$ defined above. By convention, we set $deg(0) = sdeg(0) = +\infty$.

Remark 1. We need to notice that the equalities

$$deg(fg) = deg(f) + deg(g), \qquad sdeg(fg) = sdeg(f) + sdeg(g)$$

don't hold in general for the 1-morphisms f and g, since the dot morphism labeled by v has both degree and the shifted degree equal to 1, but the square is equal to 0.

From the local relations, we get

$$QP \cong PQ \oplus \mathbf{1} \oplus \mathbf{1}\langle 1\rangle \tag{14}$$

where $\langle 1\rangle$ denotes the dimension shifting by 1 and will be explained in detail later. This can be seen by the following diagram:



which comes from the local relations.

By (10), we see that upward oriented crossings satisfy the relations of symmetric group S_n. We have a canonical homomorphism

$$\Bbbk[S_n] \longrightarrow \mathrm{End}_{'\mathcal{H}'}(P^n)$$

from the group algebra of S_n to the endomorphism ring of the n-th tensor power of P. Similarly we have a canonical homomorphism

$$\Bbbk[S_n] \longrightarrow \mathrm{End}_{'\mathcal{H}'}(Q^n).$$

For a positive integer n and a partition λ of n, let e_λ be the associated Young symmetrizer of $\Bbbk[S_n]$. The Young symmetrizers are idempotents of the group algebra $\Bbbk[S_n]$ given by

$$e_\lambda = a_\lambda b_\lambda / n_\lambda$$

where $n_\lambda = n!/\dim(V_\lambda)$ with V_λ the irreducible representation corresponding to the partition λ, the elements $a_\lambda, b_\lambda \in \Bbbk[S_n]$ are defined for a Young tableau T of the partition λ by

$$a_\lambda = \sum_{g \in L_\lambda} g, \qquad b_\lambda = \sum_{g \in L'_\lambda} sign(g) g,$$

and

$$\begin{aligned} L_\lambda &= \{g \in S_n : g \text{ preserves each row of } T\}, \\ L'_\lambda &= \{g \in S_n : g \text{ preserves each column of } T\}. \end{aligned}$$

The irreducible representations of S_n are given by $\Bbbk[S_n]e_\lambda$ for partitions λ of n. Let $e_{(n)}$ be the idempotent corresponding to the partition (n) of n, with $\Bbbk[S_n]e_{(n)}$ being the trivial representation of S_n. See [FH] for detail.

Let $'\mathcal{H}$ be the Karoubi envelope of $'\mathcal{H}'$. Then $'\mathcal{H}$ is a 2-category. The 1-morphisms P^n and Q^n splits into direct sums of (P^n, e_λ) and (Q^n, e_λ) in $'\mathcal{H}$ respectively, over partitions λ of n. Define

$$P_\lambda = (P^n, e_\lambda), \qquad Q_\lambda = (Q^n, e_\lambda), \qquad P_n = (P^n, e_{(n)}), \qquad Q_n = (Q^n, e_{(n)}).$$

Proposition 1. *There are no negative degree (respectively, shifted degree) endomorphisms of P^mQ^n, and the algebra of degree (respectively, shifted degree) zero endomorphisms of P^mQ^n is isomorphic to $\Bbbk[S_m] \otimes \Bbbk[S_n]$.*

Proof. This can be proved by looking at the string diagrams. □

Theorem 1. *In the 2-category $'\mathcal{H}$, the 1-morphisms P_n and Q_n satisfy*

$$Q_nP_m \cong \bigoplus_{k\geq 0} \bigoplus_{l=0}^{k} P_{m-k}Q_{n-k}\langle l\rangle, \tag{15}$$

$$Q_nQ_m \cong Q_mQ_n, \tag{16}$$

$$P_nP_m \cong P_mP_n. \tag{17}$$

Proof. The proof of this theorem is same as that of [CL] in the case of $\dim V = 2$. □

From Theorem 1, we get a natural homomorphism of algebras

$$\pi : \ 'H_{\mathbb{Z}} \longrightarrow K_0('\mathcal{H}) \tag{18}$$

by sending p_m to $[P_m]$ and q_n to $[Q_n]$, where $[P_m]$ and $[Q_n]$ denote the respective isomorphism classes. And we use the same notation π after coefficient extension.

Theorem 2. *π is an isomorphism of algebras.*

Proof. The proof of this theorem will be given at the end of Section 4. □

3. A Representation of the Category $'\mathcal{H}'$

We will give a construction of the 2-Category $'\mathcal{H}'$ in this section and the Fock representation of the $'\mathcal{H}$.

Let $A_1 = \mathrm{Sym}^*(V^\vee)$, which can be identified with $\Bbbk[v^\vee]$. For each positive integer n, define

$$A_n = (\mathrm{Sym}^*(V^\vee) \otimes \mathrm{Sym}^*(V^\vee) \otimes \cdots \otimes \mathrm{Sym}^*(V^\vee)) \rtimes S_n$$

where the symmetric group S_n acts by permuting the n terms in the product. The algebras inherit the natural grading from $\mathrm{Sym}^*(V^\vee)$. We denote by $\mathcal{C}_n = D(A_n\text{−gmod})$ the bounded derived category of finite dimensional, graded (left) A_n-modules. We have the maps

$$\Bbbk \xrightarrow{i} A_1 \xrightarrow{p} \Bbbk$$

with i the natural inclusion and p the projection which takes $\mathrm{Sym}^{>0}(V)$ to zero. Thus each $\Bbbk$-module is also an A_1-module and vice versa.

From the embedding $S_n = S_n \times S_1 \hookrightarrow S_{n+1}$, we have the natural inclusion $A_n \otimes A_1 \hookrightarrow A_{n+1}$. Let $P(n) := A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk)$ be an (A_{n+1}, A_n)-bimodule, and $(n)Q := (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1}$ be an (A_n, A_{n+1})-bimodule. We define the functor $\mathsf{P}(n) : D(A_n\text{−gmod}) \to D(A_{n+1}\text{−gmod})$ by

$$\mathsf{P}(n)(\cdot) := P(n) \otimes_{A_n} (\cdot), \tag{19}$$

and similarly, the functor $(n)\mathsf{Q} : D(A_{n+1}\text{−gmod}) \to D(A_n\text{−gmod})$ by

$$(n)\mathsf{Q}(\cdot) := (n)Q \otimes_{A_{n+1}} (\cdot), \tag{20}$$

with the obvious definitions of these two functors on the corresponding morphisms of the categories. In the following, we define the following natural transformations:

1. $X(1) : \mathsf{P} \to \mathsf{P}$ and $X(1) : \mathsf{Q} \to \mathsf{Q}$,
2. $X(b) : \mathsf{P} \to \mathsf{P}[1]\{-1\}$ and $X(b) : \mathsf{Q} \to \mathsf{Q}[1]\{-1\}$ for any $b \in V$,
3. $T : \mathsf{PP} \to \mathsf{PP}$, $T : \mathsf{QQ} \to \mathsf{QQ}$, $T : \mathsf{QP} \to \mathsf{PQ}$, and $T : \mathsf{PQ} \to \mathsf{QP}$,
4. $\mathrm{adj} : \mathsf{QP} \to \mathrm{id}$ and $\mathrm{adj} : \mathsf{PQ} \to \mathrm{id}[1]\{-1\}$, and
5. $\mathrm{adj} : \mathrm{id} \to \mathsf{QP}[-1]\{1\}$ and $\mathrm{adj} : \mathrm{id} \to \mathsf{PQ}$.

where $[\cdot]$ denotes the cohomological shift and $\{\cdot\}$ the grading shift. We also define the shift $\langle\cdot\rangle$ in $\mathcal{H}$ by $\langle 1\rangle := [1]\{-1\}$.

(1). $X(1) : \mathsf{P} \to \mathsf{P}$ and $X(1) : \mathsf{Q} \to \mathsf{Q}$ are the identity natural transformations in P and Q respectively.

(2). The definitions of $X(b) : \mathsf{P} \to \mathsf{P}[1]\{-1\}$ and $X(b) : \mathsf{Q} \to \mathsf{Q}[1]\{-1\}$ for any $b \in V$. Viewed as an A_1-module, a free resolution of $\Bbbk$ is:

$$\begin{array}{cccccccc} 0 & \longrightarrow & A_1 \otimes V^\vee & \xrightarrow{d} & A_1 & \longrightarrow & \Bbbk \longrightarrow & 0 \\ & & f \otimes w & \longmapsto & fw. & & & \end{array} \tag{21}$$

We define a map ϕ_b by

$$\begin{array}{rccc} \phi_b : & A_1 \otimes V^\vee & \longrightarrow & A_1\{-1\} \\ & f \otimes w & \longmapsto & \langle w, b\rangle f \end{array}$$

where $\{-1\}$ denotes the degree shifting in $A_1 = \mathrm{Sym}^*(V^\vee)$. Then we have the following commutative diagram:

$$\begin{tikzcd} 0 \arrow[r] \arrow[d] & A_1 \otimes V^\vee \arrow[r, "d"] \arrow[d, "\phi_b"] & A_1 \arrow[r] \arrow[d] & 0 \arrow[d, equal] \\ A_1 \otimes V^\vee\{-1\} \arrow[r, "-d"] & A_1\{-1\} \arrow[r] & 0 \arrow[r] & 0 \end{tikzcd}$$

which defines a morphism $X(b) : \Bbbk \longrightarrow \Bbbk[1]\{-1\}$ in the derived category $D(A_1\text{−gmod})$. This morphism then induces a morphism

$$\begin{tikzcd} X(b) : & P(n) \arrow[r] \arrow[d, equal] & P(n)[1]\{-1\} \arrow[d, equal] \\ & A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) & A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk)[1]\{-1\} \end{tikzcd}$$

Thus we get the natural transformation $X(b) : \mathsf{P} \longrightarrow \mathsf{P}[1]\{-1\}$. In the same way, we apply the following free resolution of $\Bbbk$:

$$\begin{array}{cccccccc} 0 & \longrightarrow & V^\vee \otimes A_1 & \xrightarrow{d} & A_1 & \longrightarrow & \Bbbk \longrightarrow & 0 \\ & & w \otimes f & \longmapsto & wf & & & \end{array} \tag{22}$$

and define a map ψ_b by

$$\begin{array}{rccc} \psi_b : & V^\vee \otimes A_1 & \longrightarrow & A_1\{-1\} \\ & w \otimes f & \longmapsto & -\langle w, b\rangle f \end{array}$$

Then the following commutative diagram

$$\begin{tikzcd}
 0 \arrow[r] \arrow[d] & V^\vee \otimes A_1 \arrow[r, "d"] \arrow[d, "\psi_v"] & A_1 \arrow[r] \arrow[d] & 0 \arrow[d, equal] \\
V^\vee \otimes A_1\{-1\} \arrow[r, "-d"] & A_1\{-1\} \arrow[r] & 0 \arrow[r] & 0
\end{tikzcd}$$

gives a morphism $X(b) : \Bbbk \longrightarrow \Bbbk[1]\{-1\}$ in the derived category $D(A_1{-}\mathrm{gmod})$ and induces a morphism

$$\begin{tikzcd}
X(b) : & (n)Q \arrow[r] \arrow[d, equal] & (n)Q[1]\{-1\} \arrow[d, equal] \\
 & (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} & (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1}[1]\{-1\}
\end{tikzcd}$$

Thus we get the natural transformation $X(b) : \mathsf{Q} \longrightarrow \mathsf{Q}[1]\{-1\}$.

(3). The definitions of $T : \mathsf{PP} \to \mathsf{PP}$, $T : \mathsf{QQ} \to \mathsf{QQ}$, $T : \mathsf{QP} \to \mathsf{PQ}$, and $T : \mathsf{PQ} \to \mathsf{QP}$.

- The definition of $T : \mathsf{PP} \to \mathsf{PP}$. We use the isomorphisms

$$\begin{aligned}
P(n+1) \otimes_{A_{n+1}} P(n) &\cong \big(A_{n+2} \otimes_{A_{n+1} \otimes A_1} (A_{n+1} \otimes \Bbbk)\big) \otimes_{A_{n+1}} (A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk)) \\
&\cong A_{n+2} \otimes_{A_{n+1} \otimes A_1} (A_{n+1} \otimes \Bbbk) \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) \\
&\cong A_{n+2} \otimes_{A_n \otimes A_a \otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk) \qquad (23) \\
&\cong A_{n+2} \otimes_{A_n \otimes A_2} A_n \otimes (A_2 \otimes_{A_1 \otimes A_1} (\Bbbk \otimes \Bbbk)) \qquad (24)
\end{aligned}$$

so that $P(n+1) \otimes_{A_{n+1}} P(n)$ can be identified with the expression (23) or the expression (24). We apply the map

$$\begin{aligned}
T : A_2 \otimes_{A_1 \otimes A_1} (\Bbbk \otimes \Bbbk) \quad &\longrightarrow \quad A_2 \otimes_{A_1 \otimes A_1} (\Bbbk \otimes \Bbbk) \\
a \otimes 1 \otimes 1 \quad &\longmapsto \quad a s_1 \otimes 1 \otimes 1, \qquad \text{for } a \in A_2, s_1 = (12) \in S_2,
\end{aligned}$$

where we notice that $a = ((a_1 \otimes a_2), \sigma)$ with $a_1, a_2 \in A_1, \sigma \in S_2$ and by definition $a s_1 = ((a_1 \otimes a_2), \sigma s_1)$. Therefore we get the induced map

$$\begin{aligned}
T : A_{n+2} \otimes_{A_n \otimes A_1 \otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk) \quad &\longrightarrow \quad A_{n+2} \otimes_{A_n \otimes A_1 \otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk) \\
1 \otimes (1 \otimes 1 \otimes 1) \quad &\longmapsto \quad s_{n+1} \otimes (1 \otimes 1 \otimes 1).
\end{aligned}$$

Thus we get the natural transformation $T : \mathsf{PP} \to \mathsf{PP}$.

- The definition of $T : \mathsf{QQ} \to \mathsf{QQ}$ is the same as above.
- The definition of $T : \mathsf{QP} \to \mathsf{PQ}$. Here we use the homomorphism

$$(A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) \longrightarrow A_n \otimes_{A_{n-1} \otimes A_1} (A_{n-1} \otimes \Bbbk \otimes \Bbbk) \otimes_{A_{n-1} \otimes A_1} A_n$$

of (A_n, A_n)-bimodules determined by

$$(1 \otimes 1) \otimes a \otimes (1 \otimes 1) \longmapsto \begin{cases} 0, & \text{if } a = 1; \\ 1 \otimes (1 \otimes 1 \otimes 1) \otimes 1, & \text{if } a = s_n = (n, n+1). \end{cases}$$

Then in the same way we get the mentioned natural transformation.

- The definition of $T : \mathsf{PQ} \to \mathsf{QP}$. We use the homomorphism

$$\begin{aligned}
A_n \otimes_{A_{n-1} \otimes A_1} (A_{n-1} \otimes \Bbbk \otimes \Bbbk) \otimes_{A_{n-1} \otimes A_1} A_n \quad &\longrightarrow \quad (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) \\
1 \otimes (1 \otimes 1 \otimes 1) \otimes 1 \quad &\longmapsto \quad (1 \otimes 1) \otimes s_n \otimes (1 \otimes 1)
\end{aligned}$$

to get the mentioned natural transformation.

(4). The definitions of $\mathrm{adj} : \mathsf{QP} \to \mathrm{id}$ and $\mathrm{adj} : \mathsf{PQ} \to \mathrm{id}[1]\{-1\}$.

• The definition of adj : $\mathsf{QP} \to \mathrm{id}$. It suffices to define an appropriate homomorphism of (A_n, A_n)-bimodules

$$\mathrm{adj} : (n)Q \otimes_{A_{n+1}} P(n) \longrightarrow A_n.$$

Applying the map

$$\Bbbk[S_{n+1}] \longrightarrow \Bbbk[S_n]$$

which sends 1 to 1 and $s_n = (n, n+1)$ to 0, we obtain a map $A_{n+1} \to A_n \otimes A_1$ and the induced map in the following is the required homomorphism

$$\begin{aligned} (n)Q \otimes_{A_{n+1}} P(n) = \quad & (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} \otimes_{A_{n+1}} A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) \\ \longrightarrow \quad & (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} (A_n \otimes A_1) \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk) \cong A_n. \end{aligned}$$

• The definition of adj : $\mathsf{PQ} \to \mathrm{id}[1]\{-1\}$. We need a map of (A_{n+1}, A_{n+1})-bimodules

$$P(n) \otimes_{A_n} (n)Q \to A_{n+1}[1]\{-1\}.$$

Since

$$P(n) \otimes_{A_n} (n)Q = A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1}$$

It suffices to define an appropriate map $h : \Bbbk \otimes \Bbbk \to A_1[1]\{-1\}$ of graded (A_1, A_1)-bimodules, so that we define adj as the composition of the following sequence of maps

$$\mathrm{adj} : A_{n+1} \otimes_{A_n \otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} \xrightarrow{h} A_{n+1} \otimes_{A_n \otimes A_1} A_{n+1}[1]\{-1\} \to A_{n+1}[1]\{-1\}$$

where the second map is multiplication.

In the derived category, $\Bbbk$ can be represented by the complex $0 \to A_1 \otimes V^\vee \xrightarrow{d} A_1 \to 0$ or the complex $0 \to V^\vee \otimes A_1 \xrightarrow{d} A_1 \to 0$, due to the resolutions (21) and (22), hence $\Bbbk \otimes \Bbbk$ can be represented by the complex

$$\begin{tikzcd} 0 \arrow[r] & (A_1 \otimes V^\vee) \otimes (V^\vee \otimes A_1) \arrow[r] & ((A_1 \otimes V^\vee) \otimes A_1) \oplus (A_1 \otimes (V^\vee \otimes A_1)) \arrow[r] & A_1 \otimes A_1 \arrow[r] & 0. \end{tikzcd}$$

Then we define h by the commutative diagram

$$\begin{tikzcd} 0 \arrow[r] & (A_1 \otimes V^\vee) \otimes (V^\vee \otimes A_1) \arrow[r] \arrow[d] & ((A_1 \otimes V^\vee) \otimes A_1) \oplus (A_1 \otimes (V^\vee \otimes A_1)) \arrow[r] \arrow[d] & A_1 \otimes A_1 \arrow[r] \arrow[d] & 0 \\ 0 \arrow[r] & 0 \arrow[r] & A_1\{-1\} \arrow[r] & 0 \arrow[r] & 0. \end{tikzcd}$$

where the map in the middle column is given as follows:

⋆ $(A_1 \otimes V^\vee) \otimes A_1 \to A_1$ is defined by $(f_1 \otimes \omega) \otimes f_2 \mapsto \langle \omega, v \rangle f_1 f_2$,
⋆ $A_1 \otimes (V^\vee \otimes A_1) \to A_1$ is defined by $g_1 \otimes (\omega \otimes g_2) \mapsto -\langle \omega, v \rangle g_1 g_2$.

(5). The definitions of adj : $\mathrm{id} \to \mathsf{QP}[-1]\{1\}$ **and** adj : $\mathrm{id} \to \mathsf{PQ}$.

• The definition of adj : $\mathrm{id} \to \mathsf{QP}[-1]\{1\}$. It suffices to define a map of (A_n, A_n)-bimodules

$$A_n \longrightarrow (n)Q \otimes_{A_{n+1}} P(n)[-1]\{1\}$$

which is equivalent to the map

$$A_n[1]\{-1\} \longrightarrow (A_n \otimes \Bbbk) \otimes_{A_n \otimes A_1} A_{n+1} \otimes_{A_1 \otimes A_n} (\Bbbk \otimes A_n). \tag{25}$$

We replace the first $\Bbbk$ and the second $\Bbbk$ with their isomorphic complexes by the resolutions (21) and (22). Then the map (25) comes from the following morphism of complexes in derived category

$$\begin{tikzcd}
0 \arrow[r] \arrow[d] & 0 \arrow[d] \\
0 \arrow[r] \arrow[d] & (A_n \otimes (V^\vee \otimes A_1)) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes (A_1 \otimes V^\vee)) \arrow[d] \\
A_n\{-1\} \arrow[r] \arrow[d] & \begin{array}{c}(A_n \otimes (V^\vee \otimes A_1)) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes A_1) \\ \oplus (A_n \otimes A_1) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes (A_1 \otimes V^\vee))\end{array} \arrow[d] \\
0 \arrow[r] \arrow[d] & (A_n \otimes A_1) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes A_1) = A_{n+1} \arrow[d] \\
0 \arrow[r] & 0
\end{tikzcd}$$

where $A_n\{-1\} \longrightarrow (A_n \otimes (V^\vee \otimes A_1)) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes A_1) \oplus (A_n \otimes A_1) \otimes_{A_n\otimes A_1} A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes (A_1 \otimes V^\vee))$ is uniquely determined by sending $1 \longmapsto (1 \otimes (v^\vee \otimes 1)) \otimes 1 \otimes (1 \otimes 1) - (1 \otimes 1) \otimes 1 \otimes ((1 \otimes v^\vee) \otimes 1)$. Here we need to notice that the left column has $A_n\{-1\}$ at the degree -1 position of the complex, thus represents the object $A_n[1]\{-1\}$ in the derived category. It is also easy to check that the map is well defined.

- The definition of $\mathrm{adj} : \mathrm{id} \to \mathsf{PQ}$. This functor is induced from the following map of (A_{n+1}, A_{n+1})-bimodules

$$\begin{array}{rcl}
A_{n+1} & \longrightarrow & P(n) \otimes_{A_n} (n)Q = A_{n+1} \otimes_{A_n\otimes A_1} (A_n \otimes \Bbbk \otimes \Bbbk \otimes_{A_n\otimes A_1} A_{n+1} \\
1 & \longmapsto & \sum_{i=0}^{n} s_i \cdots s_n \otimes (1 \otimes 1 \otimes 1) \otimes s_n \cdots s_i.
\end{array}$$

Here we use the convention $s_0 s_1 \cdots s_n = 1$.

Theorem 3. *The natural transformations X, T and* adj *satisfy the Heisenberg 2-relations and give a categorical Heisenberg action of ${}'\mathcal{H}$ on $\oplus_{n\geq 0} D(A_n{-}\mathrm{gmod})$.*

Proof. The proof of this theorem is very similar to that of Theorem 2 in Section 4.4 of [CL]. We will describe what needs to be proved and omit the detail. The Heisenberg 2-relations are just the local relations mentioned in Section 2, they contain the following items.

- **Adjoint relations.** The following compositions involving adjunctions are all equal to the identity 2-morphism:

(1) $\mathsf{P} \xrightarrow{I\mathrm{adj}} \mathsf{PQP}[-1]\{1\} \xrightarrow{\mathrm{adj}I} \mathsf{P}$ and $\mathsf{Q} \xrightarrow{\mathrm{adj}I} \mathsf{QPQ}[-1]\{1\} \xrightarrow{I\mathrm{adj}} \mathsf{Q}$;
(2) $\mathsf{P} \xrightarrow{\mathrm{adj}I} \mathsf{PQP} \xrightarrow{I\mathrm{adj}} \mathsf{P}$ and $\mathsf{Q} \xrightarrow{I\mathrm{adj}} \mathsf{QPQ} \xrightarrow{\mathrm{adj}I} \mathsf{Q}$.

In graphical calculus, the above relations are depicted as



That arrows are not drawn in the graph means the above equalities are true for both orientations given by (1) and (2).

- **Dots and adjunctions.** For $\forall b \in \Lambda^*(V)$, we have the following equalities of compositions of maps:

(1) $(\mathsf{QP} \xrightarrow{IX(b)} \mathsf{QP} \xrightarrow{\mathrm{adj}} \mathrm{id}) = (\mathsf{QP} \xrightarrow{X(b)I} \mathsf{QP} \xrightarrow{\mathrm{adj}} \mathrm{id})$;
(2) $(\mathsf{PQ} \xrightarrow{X(b)I} \mathsf{PQ} \xrightarrow{\mathrm{adj}} \mathrm{id}[1]\{-1\}) = (\mathsf{PQ} \xrightarrow{IX(b)} \mathsf{PQ} \xrightarrow{\mathrm{adj}} \mathrm{id}[1]\{-1\})$;

(3) $(\mathrm{id} \xrightarrow{\mathrm{adj}} \mathsf{PQ} \xrightarrow{X(b)I} \mathsf{PQ}) = (\mathrm{id} \xrightarrow{\mathrm{adj}} \mathsf{PQ} \xrightarrow{IX(b)} \mathsf{PQ})$;

(4) $(\mathrm{id} \xrightarrow{\mathrm{adj}} \mathsf{QP}[-1]\{1\} \xrightarrow{IX(b)} \mathsf{QP}[-1]\{1\}) = (\mathrm{id} \xrightarrow{\mathrm{adj}} \mathsf{QP}[-1]\{1\} \xrightarrow{X(b)I} \mathsf{QP}[-1]\{1\})$.

The above equalities can be depicted by the following:



- **Pitchfork relations.** We have the following equalities for compositions of maps:

(1) $(\mathsf{P} \xrightarrow{I\mathrm{adj}} \mathsf{PPQ} \xrightarrow{TI} \mathsf{PPQ}) = (\mathsf{P} \xrightarrow{\mathrm{adj}I} \mathsf{PQP} \xrightarrow{IT} \mathsf{PPQ})$;

(2) $(\mathsf{Q} \xrightarrow{I\mathrm{adj}} \mathsf{QPQ} \xrightarrow{TI} \mathsf{PQQ}) = (\mathsf{Q} \xrightarrow{\mathrm{adj}I} \mathsf{PQQ} \xrightarrow{IT} \mathsf{PQQ})$;

(3) $(\mathsf{PPQ} \xrightarrow{IT} \mathsf{PQP} \xrightarrow{\mathrm{adj}I} \mathsf{P}[1]\{-1\}) = (\mathsf{PPQ} \xrightarrow{TI} \mathsf{PPQ} \xrightarrow{I\mathrm{adj}} \mathsf{P}[1]\{-1\})$;

(4) $(\mathsf{PQQ} \xrightarrow{IT} \mathsf{PQQ} \xrightarrow{\mathrm{adj}I} \mathsf{Q}[1]\{-1\}) = (\mathsf{PQQ} \xrightarrow{TI} \mathsf{QPQ} \xrightarrow{I\mathrm{adj}} \mathsf{Q}[1]\{-1\})$.

The above equalities are actually rel boundary isotopy relations for 2-morphisms:



- **Dots and crossings.** The following compositions of maps with graphs containing dots are equal for $\forall b \in \Lambda^*(V)$:

(1) $(\mathsf{PP} \xrightarrow{X(b)I} \mathsf{PP} \xrightarrow{T} \mathsf{PP}) = (\mathsf{PP} \xrightarrow{T} \mathsf{PP} \xrightarrow{IX(b)} \mathsf{PP})$;

(2) $(\mathsf{PP} \xrightarrow{T} \mathsf{PP} \xrightarrow{X(b)I} \mathsf{PP}) = (\mathsf{PP} \xrightarrow{IX(b)} \mathsf{PP} \xrightarrow{T} \mathsf{PP})$.

These equalities correspond to the following graphical relations:



- **Composition of crossings relations.** The following equalities hold for composition of crossings relations:

(1) $(\mathsf{PP} \xrightarrow{T} \mathsf{PP} \xrightarrow{T} \mathsf{PP}) = (\mathsf{PP} \xrightarrow{II} \mathsf{PP})$;

(2) $(\mathsf{PPP} \xrightarrow{IT} \mathsf{PPP} \xrightarrow{TI} \mathsf{PPP} \xrightarrow{IT} \mathsf{PPP}) = (\mathsf{PPP} \xrightarrow{TI} \mathsf{PPP} \xrightarrow{IT} \mathsf{PPP} \xrightarrow{TI} \mathsf{PPP})$;

(3) $(\mathsf{PQ} \xrightarrow{T} \mathsf{QP} \xrightarrow{T} \mathsf{PQ}) = (\mathsf{PQ} \xrightarrow{II} \mathsf{PQ})$;

(4) $(\mathsf{QP} \xrightarrow{II} \mathsf{QP}) = (\mathsf{QP} \xrightarrow{T} \mathsf{PQ} \xrightarrow{T} \mathsf{QP}) + (\mathsf{QP} \xrightarrow{\mathrm{adj}} \mathrm{id} \xrightarrow{\mathrm{adj}} \mathsf{QP}[-1]\{1\} \xrightarrow{IX(v)} \mathsf{QP})$ $+ (\mathsf{QP} \xrightarrow{IX(v)} \mathsf{QP}[1]\{-1\} \xrightarrow{\mathrm{adj}} \mathrm{id}[1]\{-1\} \xrightarrow{\mathrm{adj}} \mathsf{QP})$.

These equalities correspond to the following graphical relations:



- **Counter-clockwise circles and curls.** The following equalities hold for compositions of maps:

(1) $(\mathsf{P} \xrightarrow{\mathrm{adj}I} \mathsf{QPP}[-1]\{1\} \xrightarrow{IT} \mathsf{QPP}[-1]\{1\} \xrightarrow{\mathrm{adj}I} \mathsf{P}[-1]\{1\}) = 0$;

(2) $(\mathrm{id} \xrightarrow{\text{adj}} [-1]\{1\} \xrightarrow{IX(b)} \mathsf{QP}[-1+|b|]\{1-|b|\} \xrightarrow{\text{adj}} \mathrm{id}[-1+|b|]\{1-|b|\}) = (\mathrm{id} \xrightarrow{\mathrm{tr}(b)I} \mathrm{id})$ for $\forall b \in \Lambda^*(V)$.

These equalities correspond to the following graphical relations:



$= 0$,



$= \mathrm{tr}(b)$.

□

Remark 2. In the definition (20) of the functor Q, we have $(n)\mathsf{Q} : D(A_{n+1}\text{−gmod}) \to D(A_n\text{−gmod})$ by $(n)\mathsf{Q}(\cdot) := (n)Q \otimes_{A_{n+1}} (\cdot)$. If we replace the above by a formal shifting

$$(n)\mathsf{Q}(\cdot) := (n)Q \otimes_{A_{n+1}} (\cdot)[-1/2]\{1/2\},$$

then the construction of this section goes through with a little change. Therefore we can get another categorified version of some deformed Heisenberg algebra, with the basic isomorphism (14) replaced by

$$QP \cong PQ \oplus \mathbf{1}\langle -1/2\rangle \oplus \mathbf{1}\langle 1/2\rangle.$$

The shifted degree *sdeg* in Section 2 corresponds to this construction.

4. The Grothendieck ring $K_0('\mathcal{H})$

By definition, $K_0('\mathcal{H})$ is the ring of isomorphism classes of the 1-morphisms of $'\mathcal{H}$. Let $R = \mathrm{End}('\mathcal{H}')$ be the ring of 2-morphisms, and call the string diagrams the monomials of R. Due to the basic isomorphism (14), the 1-morphisms of $'\mathcal{H}$ are of the form (M, f), where M is a finite direct sum of P^mQ^n's with m and n non-negative integers, and $f \in R$ is an idempotent. Let $R_{m,n} = \mathrm{End}(P^mQ^n)$.

Proposition 2. $K_0(R_{m,n}) \cong K_0(\Bbbk[S_m] \otimes \Bbbk[S_n])$.

Proof. Let $I = \{\varphi \in R_{m,n} : \ vdeg(\varphi) > 0\}$. Then we get a split exact sequence

$$0 \longrightarrow I \longrightarrow R_{m,n} \longrightarrow \Bbbk[S_m] \otimes \Bbbk[S_n] \longrightarrow 0.$$

Therefore we have the exact sequence for K_0 groups:

$$0 \longrightarrow K_0(I) \longrightarrow K_0(R_{m,n}) \longrightarrow K_0(\Bbbk[S_m] \otimes \Bbbk[S_n]) \longrightarrow 0. \tag{26}$$

I is a non-unital ring with an embedding $j : I \longrightarrow I_+$, where $I_+ = I \oplus \mathbb{Z}$ as abelian groups and the multiplication defined by

$$(x, m) \cdot (y, n) = (xy + my + nx, mn) \text{ for } x, y \in I \text{ and } m, n \in \mathbb{Z}.$$

For convenience, we write the element $(x, m) \in I_+$ by $x + m$. Then we have the split exact sequence

$$0 \to I \xrightarrow{j} I_+ \xrightarrow{\rho} \mathbb{Z} \to 0.$$

Then $K_0(I) = \ker(\rho_* : K_0(I_+) \longrightarrow K_0(\mathbb{Z}))$. As a unital ring, an element of $K_0(I_+)$ is some conjugacy class of an idempotent square matrix over I_+. The above approach of K_0 is refereed to [Ros]. We can view I_+ as a subring of $R_{m,n}$, thus the degree function $vdeg$ can be restricted to I_+.

For a square matrix x over I_+, let $vdeg(x)$ be the maximal $vdeg(a)$ for $a \in I_+$ taking over all non-zero entries of x, and $vdeg(0) = +\infty$. Suppose $0 \neq e \in M_t(I_+)$ is an idempotent matrix of size $t \times t$ such that $\rho_*(e) = 0$. The matrix ring can be decomposed as

$$M_t(I_+) = \bigoplus_{l=1}^{\infty} M^l$$

where M^l consists of elements of $M_t(I_+)$ with each non-zero entry having degree $l/2$. Let $M_p = \bigoplus_{l\geq p} M^l$. Then

$$M_t(I_+) = M_0 \supseteq M_1 \supseteq M_2 \supseteq \cdots, \qquad \text{and} \qquad M_p \cdot M_q \subseteq M_{p+q}.$$

We write $e = e_0 + x$ with $e_0 \in M_t(\mathbb{Z})$ and $x \in M_1$. Since e is an idempotent, we have

$$e_0^2 + e_0 x + x e_0 + x^2 = e_0 + x.$$

By comparing the degrees, we get

$$e_0^2 = e_0, \qquad \text{and} \qquad e_0 x + x e_0 + x^2 = x.$$

Since $\rho_*(e) = 0$, we have $e_0 = 0$, thus $x = x^2$ and $x \neq 0$. Hence there exists a positive integer l such that $x \in M_l \backslash M_{l+1}$, which contradicts to $x = x^2 \in M_{2l} \subseteq M_{l+1}$. Therefore we have proved that $K_0(I) = \ker(\rho) = 0$. The proposition then follows from the exact sequence (26). □

Proposition 3. *The Grothendieck ring $K_0('\mathcal{H})$ is generated by the classes $\{[P_m], [Q_n] : m, n \in \mathbb{Z}$ non-negative$\}$ as an algebra over $\Bbbk$.*

Proof. Let (M, f) be a 1-morphism of $'\mathcal{H}$. Then there exist non-negative integers m, n such that M is a direct sum of some $P^k Q^l$'s with $0 \leq k \leq m$, $0 \leq l \leq n$, and the idempotent f is a square matrix over $\mathrm{End}(P^m Q^n)$. It is well-known that $K_0(\Bbbk[S_m])$ is generated by elements of the form $[P_{i_1}] \cdot [P_{i_2}] \cdot \cdots \cdot [P_{i_t}]$ with $0 \leq i_1 + i_2 + \cdots + i_t \leq m$ as a group, hence

$$[(M,f)] = \sum c_{i_1,\cdots,i_t;j_1,\cdots,j_s} [P_{i_1}] \cdot [P_{i_2}] \cdot \cdots \cdot [P_{i_t}] \cdot [Q_{j_1}] \cdot [Q_{j_2}] \cdot \cdots \cdot [Q_{j_s}],$$

where $c_{i_1,\cdots,i_t;j_1,\cdots,j_s} \in \Bbbk$. Therefore from Proposition 2, $K_0('\mathcal{H})$ is generated by the classes $\{[P_m], [Q_n] : m, n \in \mathbb{Z}$ non-negative$\}$ as an algebra over $\Bbbk$. □

Proof of Theorem 2. At first we see that $\pi :' H \longrightarrow K_0('\mathcal{H})$ is surjective by Proposition 3. We need to prove the injectivity of π. Let $\mathcal{C}_n = D(A_n-\mathrm{gmod})$ for each integer $n \geq 0$. We have the natural map

$$j : \ '\mathcal{H} \longrightarrow \bigoplus_{n\geq 0} \mathcal{C}_n$$

which takes P_n to $(\mathsf{P}, e_{(n)})$ and Q_n to $(\mathsf{Q}, e_{(n)})$. This map induces

$$K_0(j) : \ K_0('\mathcal{H}) \longrightarrow K_0 \left(\bigoplus_{n\geq 0} \mathcal{C}_n \right),$$

hence we get the map

$$K_0('\mathcal{H}) \longrightarrow \mathrm{End}(\mathcal{F}),$$

where $\mathcal{F}$ is the Fock space representation of $'H$, as discussed in [FJW1] and [FJW2]. Consider the sequence of maps

$$'H \xrightarrow{\pi} K_0('\mathcal{H}) \xrightarrow{K_0(j)} \mathrm{End}(\mathcal{F}).$$

The composition map $K_0(j) \circ \pi$ is clearly injective, therefore π is injective. Since we already know that π is surjective, thus we get the isomorphism $\pi : \ 'H \xrightarrow{\cong} K_0('\mathcal{H})$. □

5. Categorification of MacMahon function

In this section, we will describe the coefficients of MacMahon function as the dimensions of vector spaces whose elements are string diagrams.

A 3D Young diagram can be represented by a "plane partition" [OR, ORV], i.e., a 2D array

$$\pi = (\pi_{ij})_{i,\,j=1}^{\infty} = \begin{pmatrix} \pi_{11} & \pi_{12} & \cdots \\ \pi_{21} & \pi_{22} & \cdots \\ \cdots & \cdots & \cdots \end{pmatrix} \tag{27}$$

of nonnegative integers $\pi_{ij} \in \mathbb{Z}_+$, such that$\pi_{ij} \geq \pi_{i+1,\,j}, \pi_{ij} \geq \pi_{i,\,j+1}$, where π_{ij} is the height of the stack of cubes placed at the (i, j)-th position of the plane. We will identify such a plane partition with the corresponding 3D Young diagram. The total volume of the 3D Young diagram is given by

$$\mid \pi \mid = \sum_{i,\,j=1}^{\infty} \pi_{ij}. \tag{28}$$

Given a plane partition $\pi = (\pi_{ij})_{i,j=1}^{\infty}$, the partition

$$\pi(m) = \begin{cases} (\pi_{i,\,i+m})_{i=1}^{\infty}, & \text{if } m \geq 0, \\ \\ (\pi_{j-m,\,j})_{j=1}^{\infty}, & \text{if } m < 0, \end{cases} \tag{29}$$

is the m-th diagonal slice of π. These partitions $\{\pi(m)\}_{m=-\infty}^{\infty}$ represent a sequence of 2D Young diagrams that are literally obtained by slicing the 3D Young diagrams diagonally.

The diagonal slices are not arbitrary but satisfy the condition

$$\cdots \prec \pi(-2) \prec \pi(-1) \prec \pi(0) \succ \pi(1) \succ \pi(2) \succ \cdots \tag{30}$$

where $\succ$ denotes interlacing relation. We say two partitions λ and μ of 2D Young diagram satisfy the interlacing relation

$$\lambda = (\lambda_1,\, \lambda_2,\, \lambda_3, \cdots) \succ \mu = (\mu_1,\, \mu_2,\, \mu_3, \cdots)$$

if

$$\lambda_1 \geq \mu_1 \geq \lambda_2 \geq \mu_2 \geq \lambda_3 \geq \cdots .$$

Let $P(N)$ denotes the number of partition π of 3D Young diagram with $\mid \pi \mid = N$. The generating function of these numbers was studied by MacMahon[B] and showed to be given by the MacMahon function:

$$\sum_{N=0}^{\infty} P(N) q^N = \Pi_{n=1}^{\infty} (1 - q^n)^{-n}. \tag{31}$$

In statistical mechanics, this generating function becomes the partition function

$$Z = \sum_{N=0}^{\infty} P(N) q^N = \sum_{\pi} q^{|\pi|} \tag{32}$$

of a canonical ensemble of plane partitions, in which each plane partition π has an energy proportional to the volume $\mid \pi \mid$.

Consider the operators

$$\Gamma_{\pm}(z) = exp\,(\sum_{n>0} \frac{a_{\pm n} z^n}{n}) \tag{33}$$

where a_n denotes the modes of the fermionic current $\phi^* \phi$ in Equation (3). The operators $\Gamma_{\pm}(z)$ can be identified with annihilation and creation parts of the bosonic vertex operator $e^{\phi(z)}$. One can get the following commutation relation

$$\Gamma_+(z)\Gamma_-(w) = \frac{1}{1-zw}\Gamma_-(w)\Gamma_+(z). \tag{34}$$

The vertex operators $\Gamma_\pm(z)$ act on the aforementioned orthonormal bases $|\lambda\rangle$ and $\langle\lambda|$ as

$$\begin{aligned} \langle\lambda|\Gamma_+(z) &= \sum_{\mu\succ\lambda}\langle\mu|z^{(|\mu|-|\lambda|)} \\ \Gamma_-(z)|\lambda\rangle &= \sum_{\mu\succ\lambda} z^{(|\mu|-|\lambda|)}|\mu\rangle. \end{aligned} \tag{35}$$

By identification (35) of the transfer matrix, we have

$$Z = \langle vac|\cdots\Gamma_+(q^{\frac{3}{2}})\Gamma_+(q^{\frac{1}{2}})\Gamma_-(q^{\frac{1}{2}})\Gamma_-(q^{\frac{3}{2}})\cdots|vac\rangle \tag{36}$$

Let

$$\widetilde{\Gamma}_-(z) = \sum_{n=0}^{\infty} P_n z^n, \tag{37}$$

$$\widetilde{\Gamma}_+(z) = \sum_{n=0}^{\infty} Q_n z^n, \tag{38}$$

where P_n, Q_n are 1-morphisms in 2-category $'\mathcal{H}$ described in Section 2, and z an indeterminate.

Proposition 4. *The following holds:*

$$\widetilde{\Gamma}_-(z)V_\lambda = \sum_{n=0}^{\infty} P_n z^n V_\lambda = \sum_{n=0}^{\infty} \sum_{\substack{\mu\succ\lambda,\\ |\mu|=|\lambda|+n}} V_\mu\, z^n \tag{39}$$

where V_λ denotes the irreducible representation of the symmetric group $S_{|\lambda|}$ associated to the partition λ.

Proof. This result should be found in principle in [FH] and [M]. As explained in Section 3, the representation of P is the induction functor from $\Bbbk[S_d]-$mod $\to \Bbbk[S_{d+1}]-$mod, which takes V_λ to $\mathrm{Ind}(V_\lambda)$ for a partition λ of d, with $\mathrm{Ind}(V_\lambda)$ equal to the direct sum of all V_μ and μ satisfying $|\mu| = 1+|\lambda|$, $\lambda \subset \mu$. The representation $P_n V_\lambda$ is isomorphic to $V_\lambda \otimes V_{(n)}$, which can be decomposed as

$$V_\lambda \otimes V_{(n)} = \sum_{\substack{\mu\succ\lambda,\\ |\mu|=|\lambda|+n}} V_\nu,$$

where $V_{(n)}$ denotes the trivial representation of S_n. □

We define the partition function $\widetilde{Z}$ of the functors P_n, Q_n for $n \geq 0$ by

$$\widetilde{Z} = \langle vac|\cdots\widetilde{\Gamma}_+(q^{\frac{3}{2}})\widetilde{\Gamma}_+(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\cdots|vac\rangle.$$

Proposition 5. *The partition function $\widetilde{Z}$ is*

$$\widetilde{Z} = \prod_{n=1}^{\infty}\left(\frac{1}{1-q^n}\right)^n \cdot \prod_{n=1}^{\infty}\left(\frac{1}{1-tq^n}\right)^n.$$

Proof. From Equations (15)–(17) of Theorem 1, we get the commutation relation

$$\widetilde{\Gamma}_+(z)\widetilde{\Gamma}_-(w) = \widetilde{\Gamma}_-(w)\widetilde{\Gamma}_+(z)\frac{1}{1-zw}\frac{1}{1-tzw},$$

hence

$$\widetilde{\Gamma}_+(q^{k/2})\widetilde{\Gamma}_-(q^{l/2}) = \widetilde{\Gamma}_-(q^{l/2})\widetilde{\Gamma}_+(q^{k/2})\frac{1}{1-q^{(k+l)/2}}\frac{1}{1-tq^{(k+l)/2}}$$

for odd positive integers k and l, which gives rise to the formula for the partition function $\widetilde{Z}$. ☐

Remark 3. If we apply the functors P_n and Q_n in the categorificatin of the Heisenberg algebra as explained in Remark 2, then the partition function $\widetilde{Z}$ equals

$$\prod_{n=1}^{\infty}\left(\frac{1}{1-t^{-1/2}q^n}\right)^n \cdot \prod_{n=1}^{\infty}\left(\frac{1}{1-t^{1/2}q^n}\right)^n.$$

The classical MacMahon function Z is also seen to be related to the graded vector space of 2-morphisms of $'\mathcal{H}$. Let $V_0 = \Bbbk \cdot 1$ be the 1-dimensional vector space over k. It is clear that $\mathrm{End}(\widetilde{\Gamma}_-(q^{\frac{1}{2}})$ $\widetilde{\Gamma}_-(q^{\frac{3}{2}})$ $\widetilde{\Gamma}_-(q^{\frac{5}{2}})\cdots)$ is a graded vector space. We write

$$\mathrm{End}(\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\widetilde{\Gamma}_-(q^{\frac{5}{2}})\cdots) = \bigoplus_{i=0}^{\infty} W_i(q)$$

where elements of $W_i(q)$ have degree i and the grading depends on q. Then we define

$$\mathscr{Z}(q,t) = \mathrm{gdim}\,\mathrm{End}(\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\widetilde{\Gamma}_-(q^{\frac{5}{2}})\cdots) = \sum_{i=0}^{\infty} \dim W_i(q)t^i.$$

Proposition 6. $\mathscr{Z}(q,0) = W_0(q) = Z$.

Proof. By continuing composing with the adjoint map  , we get a degree preserving isomorphism

$$\mathrm{End}(\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\widetilde{\Gamma}_-(q^{\frac{5}{2}})\cdots) \cong \mathrm{Hom}(V_0\cdots\widetilde{\Gamma}_+(q^{\frac{3}{2}})\widetilde{\Gamma}_+(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\cdots, V_0).$$

By Equations (15)–(17) of Theorem 1, we can exchange the composition order of those transfer matrice, and need only count the number of terms of $\mathbf{1}$ in the expansions of those $Q_{m_1}\cdots Q_{m_s}\cdot P_{n_1}\cdots P_{n_t}$ in

$$\cdots\widetilde{\Gamma}_+(q^{\frac{5}{2}})\widetilde{\Gamma}_+(q^{\frac{3}{2}})\widetilde{\Gamma}_+(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{1}{2}})\widetilde{\Gamma}_-(q^{\frac{3}{2}})\widetilde{\Gamma}_-(q^{\frac{5}{2}})\cdots.$$

Therefore we get $\mathscr{Z}(q,0) = W_0(q) = Z$. ☐

The authors of this paper don't know how to calculate the function $\mathscr{Z}(q,t)$ or relate it to other topics. It is speculate that the categorified version of the Heisenberg algebra should play some role in the quantum field theory, invariant theory, etc.. We hope further studies would reveal something interesting in the future.

References

[B] D.M. Bressoud, Proofs and confirmations, The story of the alternating sign matrix conjecture, Cambridge University Press, 1999.

[CL] S. Cautis, A. Licata, Heisenberg categorification and Hibert schemes. arXiv: math.QA/1009.5147.

[FH] W. Fulton, J. Harris, Representation theory, A first course. Springer-Verlag, New York, 1991.

[FJW1] I. Frenkel, N. Jing, and W. Wang, Vertex Representations via finite groups and the McKay corrrespondence, *Int. Math. Res. Not.* (2000), no. 4, 195–222.

[FJW2] I. Frenkel, N. Jing, and W. Wang, Quantum vertex representations via finite groups and the McKay correspondence, *Comm. Math. Phys.* **211** (2000), no. 2, 365–393.

[K] M. Khovanov, Heisenberg algebra and a graphical calculus; arXiv:1009.3295.

[M] I. G. Macdonald, Symmetric functions and Hall polynomials. Oxford Mathematical Monographs, Clarendon Press, Oxford, 1979.

[MJD] T. Miwa, M. Jimbo and E. Date, Solitons: Differential equations, symmetries and infinite dimensional algebras, Cambridge University Press (2000).
[OR] A. Okounkov and N. Reshetikhin, Correlation function of Schur Process with application to local geometry of a random 3-Dimensional young diagram, J. Amer. Math. Soc. 16, (2003), 581C603.
[ORV] A. Okounkov, N. Reshetikhin and C. Vafa, Quantum Calabi-Yau and classical crystals, in: P. Etingof, V. Retakh and I.M. Singer (eds.), The unity of mathematics, Progr. Math. 244, Birkhauser, 2006, pp. 597C618.
[Ros] J. Rosenberg, Algebraic K-Theory and Its Applications, GTM 147, Springer-Verlag, New York, 1994.

Na Wang, College of mathematics and information sciences, Henan University, Kaifeng, 475001, China,
E-mail address: `wangnawanda@163.com`

Zhixi Wang, School of mathematical science, Capital Normal University, Beijing, 100048, China,
E-mail address: `wangzhx@mail.cnu.edu.cn`

Ke Wu, School of mathematical science, Capital Normal University, Beijing, 100048, China,
E-mail address: `wuke@mail.cnu.edu.cn`

Jie Yang, School of mathematical science, Capital Normal University, Beijing, 100048, China,
E-mail address: `yangjie@cnu.edu.cn`

Zifeng Yang, School of mathematical science, Capital Normal University, Beijing, 100048, China
E-mail address: `yangzf@mail.cnu.edu.cn`